

Dibaryon Systems in Chiral $SU(3)$ Quark Model¹

Q.B.Li^a, P.N.Shen^{b,a,c,d}, Z.Y.Zhang^a, Y.W.Yu^a

- a.* Institute of High Energy Physics, Academia Sinica, P.O.Box 918(4),
Beijing 100039, China
- b.* China Center of Advanced Science and Technology (World
Laboratory), P.O.Box 8730, Beijing 100080, China
- c.* Institute of Theoretical Physics, Academia Sinica,
P.O.Box 2735, Beijing 100080, China
- d.* Center of Theoretical Nuclear Physics, National Lab of Heavy
Ion Accelerator, Lanzhou 730000, China

Abstract

The possible candidates of S -wave dibaryons with various strange numbers are studied under the chiral $SU(3)$ quark model. It is shown that there are three types of baryon-baryon bound states. The states of the first type are called deuteron-like states. If chiral fields can provide enough attraction between interacting baryons, these systems, such as $[\Xi\Omega - \Xi^*\Omega]_{(1,1/2)}$, $[\Xi\Xi]_{(0,1)}$, $[N\Omega]_{(2,1/2)}$ would be weakly bound. The states of the second type such as $[\Sigma^*\Delta]_{(0,5/2)}$, $[\Sigma^*\Delta]_{(3,1/2)}$, $[\Delta\Delta]_{(0,3)}$ and $[\Delta\Delta]_{(3,0)}$ are named as $\Delta\Delta$ -like states. Due to the highly symmetric character in orbital space, these systems could be relatively deeply bound, but the strong decay modes of composed baryons cause the widths of the states much broader. The states of the third type are entitled as $\Omega\Omega$ -like states. Due to the same symmetry character shown in the systems of the second type and the only weak decay mode of composed baryons, for instance in $[\Omega\Omega]_{(0,0)}$, or at most one strong decay mode of composed baryons, for example in $[\Xi^*\Omega]_{(0,1/2)}$, these states are deeply bound states with narrow widths. The states of latter two types are most interesting new dibaryon states and should be carefully investigated both theoretically and experimentally.

¹This work was partly supported by the National Natural Science Foundation of China (NSFC) and the Chinese Academy of Sciences

1. Introduction.

Dibaryon as a six-quark system has shown its special place in the investigation of medium-energy physics since Jaffe published his prediction of H particle in 1977 [1]. As is commonly believed, to study a quark system, the Quantum Chromodynamics (QCD), which governs the strong interaction among quarks and gluons, should be employed as an underline theory although its non-perturbative behavior is still not quite clear and cannot exactly be solved up to now. Jaffe studied the color-magnetic interaction (CMI) of the one-gluon-exchange (OGE) potential in the multi-quark system and found that CMI shows attractive character in the H particle case. This character compels six quarks staying in a smaller volume, say less than $0.85fm$ in radius. Thus dibaryon study could provide more information about the short-range behavior of QCD , and the existence of dibaryon can directly supply the evidence of the quark-gluon degrees of freedom in hadrons and hadronic systems. However, the reason for forming the baryon-baryon bound state presents great complexity. The nonperturbative QCD ($NPQCD$) effect may seriously affect the properties of the dibaryon due to its finite size. The CMI and the interaction describing the action from physical vacuum should be co-responsible for the binding energy of the system. The symmetry structure of the system may also play an important role there. In a word, the character of CMI is no longer dominant. Sometimes, meson clouds may provide predominant effects. Exploring dibaryon may enable us to investigate the short- and medium-range $NPQCD$ effects and to find out a practical way to properly treat them.

In order to reliably study dibaryon, a model that can describe most of short- and medium-range $NPQCD$ effects should be employed. In other word, the model should have predictive power. It should at least contain two-fold requirements: By using this model the ground state properties of baryons should well be fitted, and the experimental data of the nucleon-nucleon ($N - N$) scattering and, especially, the empirical data of the nucleon-hyperon ($N - Y$) scattering and reaction can reasonably be reproduced in the dynamical calculation. When extending this model to dibaryon investigation, no additional parameters are required. There are lots of models such as MIT bag model [2], the constituent quark models of various kinds [3, 7, 8], Skyrme model [9], etc. Among them, the chiral $SU(3)$ quark model [7, 8] is one of the models which

satisfy requested conditions. In terms of this model, investigating and further systematically analyzing possible bound six-quark systems become significant and essential.

Since 70's, dibaryon has been intensively studied. The most interesting dibaryons have been studied are the following: H particle has been theoretically and experimentally investigated for years. The theoretically predicted mass is in a large range [1, 3, 4, 5, 6, 15, 26], say from 2GeV to 2.4GeV . The most believed theoretical prediction is around the $\Lambda\Lambda$ threshold, namely around 2.232GeV [4, 5]. However, this particle still has not been found in the experiment yet. The most recent data showed that the lower limit of the H particle mass is about 2.22GeV [13]. Except the H particle, possible bound baryon-baryon systems in the non-strangeness sector were also investigated. d^* is one of them. There were number of theoretical predictions by using various models, such as the non-relativistic boson-exchange model [17], the quark cluster model [4], the quark-delocation model [19], the chiral $SU(3)$ quark model [20] and etc.. The predicted masses also spread in a wide range. All the predictions are below the threshold of the $\Delta\Delta$ channel of 2.464GeV , and most of them are above the threshold of the strong decay channel, $NN\pi\pi$, of 2.154GeV . $d'(J^P = 0^-, T = 0)$ is another interesting particle. In the experiments of double-charge-exchange reactions, it was found that when the energy of the incident pion is 50MeV , there exists a resonance with the mass of 2.065GeV and the width of 0.5MeV in the processes with a variety of targets [22]. To explain that phenomenon, one proposed d' . Although there were many theoretical attempts [23], the theoretical result is still away from the expected value. Whether this phenomenon indicates d' is still under discussion. Up to now, these three interesting candidates of dibaryons are still not found or confirmed by experiments. It seems that one should go beyond these candidates and should search the possible candidates in a wider region, especially the systems with multi-strangeness, in terms of a more reliable model such as chiral $SU(3)$ quark model. According to this idea, Yu et al. analyzed the six-quark system with a simple six-quark cluster configuration [16]. Later, by employing the chiral $SU(3)$ quark model, Zhang, Yu et al. studied $\Omega\Omega(S = 0, T = 0)$ and $\Xi\Omega(S = 1, T = 1/2)$ [16, 14], and Li and Shen explored $\Xi^*\Omega(S = 0, T = 1/2)$ and $\Xi\Omega - \Xi^*\Omega(S = 1, T = 1/2)$ [24]. In this paper, we would present a sys-

tematic study of possible candidates of S -wave baryon-baryon bound states in this model.

The paper is arranged in the following way: The chiral $SU(3)$ quark model is briefly introduced in Sect.2. In Sect.3, the results calculated by this model are given, and the symmetry characters of the system concerned are discussed. The effects of chiral-quark field induced interactions on the binding behaviors of systems are detailed analyzed in Sect.4. In Sect.5, the model parameter dependence of the predicted binding energy is further studied. Finally, in Sect.6, the concluding remark is drawn.

2. Brief introduction of chiral $SU(3)$ quark model.

Following Georgi's idea [10], the quark-chiral $SU(3)$ field interaction can be written as

$$\mathcal{L}_I = -g_{ch}(\bar{\psi}_L \Sigma \psi_R - \bar{\psi}_R \Sigma^+ \psi_L) , \quad (1)$$

with g_{ch} being the quark-chiral field coupling constant, ψ_L and ψ_R being the quark-left and right spinors, respectively, and

$$\Sigma = \exp[i\pi_a \lambda_a / f], \quad a = 1, 2, \dots, 8 . \quad (2)$$

where π_a is the Goldstone boson field and λ_a the Gell Mann matrix of the flavor $SU(3)$ group. Generalizing the linear realization of Σ in the $SU(2)$ case to the $SU(3)$ case, one obtains

$$\Sigma = \sum_{a=0}^8 \sigma_a \lambda_a - i \sum_{a=0}^8 \pi_a \lambda_a \gamma_5 , \quad (3)$$

and the interaction Lagrangian

$$\mathcal{L}_I = -g_{ch} \bar{\psi} \left(\sum_{a=0}^8 \sigma_a \lambda_a + i \sum_{a=0}^8 \pi_a \lambda_a \gamma_5 \right) \psi , \quad (4)$$

where λ_0 is a unitary matrix, $\sigma_0, \dots, \sigma_8$ the scalar nonet fields and $\pi_0, \dots, \pi_8$ the pseudo-scalar nonet fields. Clearly, $\mathcal{L}_I$ is invariant under the infinitesimal chiral $SU(3)_L \times SU(3)_R$ transformation. Consequently, one can write the interactive Hamiltonian as

$$H_{ch} = g_{ch} F(q^2) \bar{\psi} \left(\sum_{a=0}^8 \sigma_a \lambda_a + i \sum_{a=0}^8 \pi_a \lambda_a \gamma_5 \right) \psi . \quad (5)$$

Here we have inserted a form factor $F(q^2)$ to describe the chiral field structure [11]. As usual, $F(q^2)$ is taken as

$$F(q^2) = \left(\frac{\Lambda_{CSB}^2}{\Lambda_{CSB}^2 + q^2} \right)^{1/2}, \quad (6)$$

and the cut-off mass Λ_{CSB} indicates the chiral symmetry breaking scale [11].

Then, the $SU(3)$ chiral-field-induced quark-quark potentials can be derived in the following :

$$\begin{aligned} V_{\sigma_a}(\vec{r}_{ij}) &= -C(g_{ch}, m_{\sigma_a}, \Lambda_{CSB}) X_1(m_{\sigma_a}, \Lambda_{CSB}, r_{ij}) (\lambda_a(i) \lambda_a(j)) \\ &+ V_{\sigma_a}^{\vec{L}\vec{S}}(\vec{r}_{ij}), \end{aligned} \quad (7)$$

$$\begin{aligned} V_{\pi_a}(\vec{r}_{ij}) &= C(g_{ch}, m_{\pi_a}, \Lambda_{CSB}) \frac{m_{\pi_a}^2}{12m_{qi}m_{qj}} [X_2(m_{\pi_a}, \Lambda_{CSB}, r_{ij}) (\vec{\sigma}_i \cdot \vec{\sigma}_j) \\ &+ \left(H(m_{\pi_a} r_{ij}) - \left(\frac{\Lambda_{CSB}}{m_{\pi_a}} \right)^3 H(\Lambda_{CSB} r_{ij}) \right) S_{ij}] (\lambda_a(i) \lambda_a(j)) \quad , \end{aligned} \quad (8)$$

with

$$\begin{aligned} V_{\sigma_a}^{\vec{L}\vec{S}}(\vec{r}_{ij}) &= -C(g_{ch}, m_{\sigma_a}, \Lambda_{CSB}) \frac{m_{\sigma_a}^2}{4m_{qi}m_{qj}} [G(m_{\sigma_a} r_{ij}) \\ &- \left(\frac{\Lambda_{CSB}}{m_{\sigma_a}} \right)^3 G(\Lambda_{CSB} r_{ij})] (\vec{L} \cdot (\vec{\sigma}_i + \vec{\sigma}_j)) (\lambda_a(i) \lambda_a(j)) \quad , \end{aligned} \quad (9)$$

$$C(g_{ch}, m, \Lambda) = \frac{g_{ch}^2}{4\pi} \frac{\Lambda^2}{\Lambda^2 - m^2} m \quad , \quad (10)$$

$$S_{ij} = 3(\vec{\sigma}_i \cdot \hat{r})(\vec{\sigma}_j \cdot \hat{r}) - (\vec{\sigma}_i \cdot \vec{\sigma}_j) \quad , \quad (11)$$

and m_{σ_a} being the mass of the scalar meson and m_{π_a} the mass of the pseudo-scalar meson. The explicit forms of functions X_1 , X_2 , H and G can be found in [7].

As mentioned in [7], the interactions induced by chiral fields describe the NPQCD effect in the low-momentum medium-distance range, which is very important in explaining the short- and medium-range forces between two baryons. To study the baryon structure and baryon-baryon dynamics, one still needs an effective one-gluon-exchange interaction V_{ij}^{OGE} which dominates the short-range perturbative QCD behavior and a confinement potential V_{ij}^{conf} which provides the NPQCD effect in the long distance and confines three quarks to a baryon. Then, the total Hamiltonian of the six-quark system can be written

as

$$H = \sum_i T_i - T_G + \sum_{i < j} V_{ij} \quad , \quad (12)$$

with

$$V_{ij} = V_{ij}^{OGE} + V_{ij}^{conf} + V_{ij}^{ch} \quad , \quad (13)$$

where

$$\begin{aligned} V_{ij}^{OGE} &= \frac{1}{4} g_i g_j (\lambda_i^c \cdot \lambda_j^c) \left\{ \frac{1}{r_{ij}} - \frac{\pi}{2} \delta(\vec{r}_{ij}) \left(\frac{1}{m_{qi}^2} + \frac{1}{m_{qj}^2} \right) \right. \\ &\quad \left. + \frac{4}{3} \frac{1}{m_{qi} m_{qj}} (\vec{\sigma}_i \cdot \vec{\sigma}_j) \right\} - \frac{1}{4 m_{qi} m_{qj} r_{ij}^3} S_{ij} \} + V_{OGE}^{\vec{\ell}, \vec{s}} \quad , \end{aligned} \quad (14)$$

$$V_{OGE}^{\vec{\ell}, \vec{s}} = -\frac{1}{16} g_i g_j (\lambda_i^c \cdot \lambda_j^c) \frac{3}{m_{qi} m_{qj}} \frac{1}{r_{ij}^3} \vec{L} \cdot (\vec{\sigma}_i + \vec{\sigma}_j) \quad , \quad (15)$$

$$V_{ij}^{conf} = -a_{ij}^c (\lambda_i^c \cdot \lambda_j^c) r_{ij}^2 - a_{ij}^{c0} (\lambda_i^c \cdot \lambda_j^c) \quad , \quad (16)$$

and

$$V_{ij}^{ch} = \sum_{a=0}^8 V_{\sigma_a}(\vec{r}_{ij}) + \sum_{a=0}^8 V_{\pi_a}(\vec{r}_{ij}) \quad . \quad (17)$$

The model parameter should be fixed before calculation. The coupling constant g_{ch} is fixed by

$$\frac{g_{ch}^2}{4\pi} = \left(\frac{3}{5}\right)^2 \frac{g_{NN\pi}^2}{4\pi} \frac{m_q^2}{M_N^2} \quad , \quad (18)$$

and $g_{NN\pi}^2/4\pi$ is taken to be the empirical value of about 14. The masses of the pseudo-scalar meson $m_\pi, m_\eta, m_{\eta'}$ and m_K can be chosen as the masses of the real π, η, η' and K , and the mass of the scalar meson σ_0 can be taken as $m_{\sigma_0} \simeq 625 \text{ MeV}$, according to the relation [12]

$$m_{\sigma_0}^2 = (2m_q)^2 + m_\pi^2 \quad . \quad (19)$$

In our previous investigation [7], it was found the $N-N$ and $N-Y$ scatterings are not sensitive to the masses of strange chiral fields. In order to reduce the numbers of adjustable parameters, m_{σ_a} ($a = 1, \dots, 8$) are also taken to be the mass of η' . The cut-off mass Λ_{CSB} for various chiral fields is taken to be

$$\Lambda_{CSB} = \begin{cases} 4.2 fm^{-1}, & \text{for } \pi, K \text{ and } \sigma_0 \\ 5.0 fm^{-1}, & \text{for } \eta, \eta', \sigma', \kappa \text{ and } \epsilon. \end{cases} \quad (20)$$

This set of model parameters is called Model I (Set I) which was frequently used in our pervious investigations [7, 8, 5]. Because the systems studied in this paper mostly comprise strange quarks, the strange chiral clouds surrounding the baryons become influential. In order to see this effect, we increase the masses and corresponding cut-masses of κ and ϵ to the values of 1.4GeV and 1.5GeV , respectively, which are close to the masses of real mesons with the same quantum numbers [25]. This set of parameters is called Model I (Set II). When the values of $m_u, m_s, b_u, g_{ch}, m_{\pi_a}, m_{\sigma_a}$ and Λ_{CSB} are fixed, the one gluon exchange coupling constants g_u and g_s can be determined by mass splittings between Δ and N , and Σ and Λ , respectively, the confinement strenghts a_{uu}^c , a_{us}^c and a_{ss}^c are fixed by the stability conditions of N , Λ and Ξ , respectively, and the zero point energies a_{uu}^{c0} , a_{us}^{c0} and a_{ss}^{c0} are fixed by the masses of N , Λ and $\overline{\Sigma} + \overline{\Omega}$, respectively.

As is mentioned above, to predict the dibaryon structure, the model should be able to reproduce the data of the N-N and Y-N scatterings reasonably. The detailed comparison of the theoretical scattering results and the empirical data can be found in [7]. Here, we only show a typical plot, the cross section of the $\Lambda - p$ process in Fig.1. In this figure, the solid and dashed curves represent the results with Sets I and II, respectively. It is shown that both curves are consistent with the experimental data. After confirming the model, we use the same set of parameters to study the dibaryon system.

All the model parameters in Sets I and II are tabulated in Table 1, respectively.

Table 1. Model parameters

	Set I	Set II
$m_u(MeV)$	313	313
$m_s(MeV)$	470	470
$b_u(fm)$	0.505	0.505
g_u	0.936	0.936
g_s	0.924	0.781
$a_{uu}^c(MeV/fm^2)/a_{uu}^{c0}(MeV)$	54.3/-47.7	55.0/-48.9
$a_{us}^c(MeV/fm^2)/a_{us}^{c0}(MeV)$	65.8/-41.7	66.5/-50.6
$a_{ss}^c(MeV/fm^2)/a_{ss}^{c0}(MeV)$	103.0/-50.6	115.4/-73.7
$m_\pi(fm^{-1})/\Lambda_\pi(fm^{-1})$	0.7/4.2	0.7/4.2
$m_K(fm^{-1})/\Lambda_K(fm^{-1})$	2.51/4.2	2.51/4.2
$m_\eta(fm^{-1})/\Lambda_\eta(fm^{-1})$	2.78/5.0	2.78/5.0
$m_{\eta'}(fm^{-1})/\Lambda_{\eta'}(fm^{-1})$	4.85/5.0	4.85/5.0
$m_{\sigma_0}(fm^{-1})/\Lambda_{\sigma_0}(fm^{-1})$	3.17/4.2	3.17/4.2
$m_{\sigma'}(fm^{-1})/\Lambda_{\sigma'}(fm^{-1})$	4.85/5.0	4.85/5.0
$m_\kappa(fm^{-1})/\Lambda_\kappa(fm^{-1})$	4.85/5.0	7.09/7.6
$m_\epsilon(fm^{-1})/\Lambda_\epsilon(fm^{-1})$	4.85/5.0	7.09/7.6

The binding energy of the baryon-baryon system is dynamically solved by using the Resonating Group Method (*RGM*). In this method, the trial wave function of the six-quark system can be written as

$$\Psi = \mathcal{A}[\hat{\phi}_A(\xi_1, \xi_2)\hat{\phi}_B(\xi_4, \xi_5)\chi_{rel}(\vec{R})\chi_{CM}(\vec{R}_{CM})]_{ST}, \quad (21)$$

where $\phi_{A(B)}$ denotes the antisymmetrized wave function of the baryon cluster A(B), $\chi_{rel}(\vec{R})$ the trial wave function of the relative motion between interacting clusters A and B , $\chi_{CM}(\vec{R}_{CM})$ the wave function of the motion of the total center of mass, and ξ_i the Jacobi coordinate with $i = 1$ and 2 for cluster A and $i = 4$ and 5 for cluster B , respectively. The symbol $\mathcal{A}$ describes the operation of the antisymmetrization between quarks in two interacting clusters. This operator can be read as

$$\mathcal{A} = N \sum_P \epsilon_P P,$$

where P is an operator which permutes quarks of cluster A and quarks of cluster B , $\epsilon_P = 1(-1)$ when P is an even(odd) permutation and N is the normalization factor. Considering the permutation symmetry, $\mathcal{A}$ can also be written as

$$\mathcal{A} = N'(1 - \sum_{\substack{i \in A \\ j \in B}} P_{ij}^{osfc}), \quad (22)$$

where P_{ij}^{osfc} denotes the permutation operation carried out between the i -th and j -th quarks in the *orbit*, *spin*, *flavor* and *color* spaces, simultaneously, and again N' represents the normalization constant. Calculating the expectation value of the Hamiltonian operator on the trial wave function in which the unknown $\chi_{rel}(\vec{R})$ is expanded in terms of well-known bases, one deduces a secular equation

$$\sum_{j=1}^n (H_{ij} - EN_{ij})c_j = 0, \quad (i = 1, 2, \dots, n), \quad (23)$$

with H_{ij} and N_{ij} being the Hamiltonian and normalization matrix elements, respectively, E the eigenvalue and c_i the corresponding eigenfunction, namely the expansion coefficients of $\chi_{rel}(\vec{R})$. Solving this equation for c_j , one obtains the binding energy and the corresponding wave function of the six-quark system. The detailed method can be found in our previous paper [24, 5].

3. Symmetry character discussion.

As a comprehensive survey, there are two crucial physical factors which resolve whether a two-baryon system is bound. One is the symmetry property of the system, namely the characteristics of quark exchanges between baryons, and the other is the interaction including both the direct and exchange components between quarks, and eventually between baryons. In this section, we analyze the symmetry property of the system according to the character of the matrix element P_{36}^{sfc} , where the superscript *sfc* denotes the P operator acts within the *spin* – *flavor* – *color* space only and the subscript 36 represents the exchange operation is between the 3-rd and 6-th quarks. Then, in section 4, we discuss the effects of various chiral-quark field interactions by employing several models.

As is pointed out in Ref.[21], the matrix element $\langle \mathcal{A}^{sfc} \rangle$ is an important measure of the action of the Pauli principle in the two-baryon state. This measure specifies the symmetry character of the state of the system. According to the symmetry characters of systems, namely the mentioned matrix elements, the two-baryon systems concerned can generally be divided into three classes. In the first class, $9\langle P_{36}^{sfc} \rangle \sim 1$, namely, $\langle \mathcal{A}^{sfc} \rangle \sim 0$. The Pauli blocking effect between interacting baryons are incredibly serious so that the two-baryon S -wave state with $[6]_r$, the $[6]$ symmetry in the orbital space, is almost a for-

bidden state. Namely, it is very hard to form a bound state. The state in the second class has the property of $9\langle P_{36}^{sfc} \rangle \sim 0$, or $\langle \mathcal{A}^{sfc} \rangle \sim 1$. In this class, the Pauli blocking effect between interacting baryons are very small so that the exchange effect between quarks which are located in different baryons, respectively, becomes negligible and these two baryons are relatively independent with each other. As a result, the meson-exchange effect may play a dominant role in binding. If the inter-baryon interaction shows an attractive feature with a large enough strength, the system would be bound. This kind of system may also be deduced in terms of a model in the baryon-meson degrees of freedom. The state in the third class possesses a feature of $9\langle P_{36}^{sfc} \rangle \sim -1$, or $\langle \mathcal{A}^{sfc} \rangle \sim 2$. The inter-baryon quark-exchange feature of this kind would be enormously beneficial to form a state with the $[6]_r$ symmetry [21]. If the inter-baryon interaction demonstrates the attractive character with certain strength, it is possible to form not only a bound state, but also a dibaryon with a relative smaller size in radius. We present the resultant binding energies and the corresponding root-mean-square radii (RMS) of the S -states which have various strange number $\mathcal{S}$ and belong to the second and third classes in the following subsections.

3.1. The systems in which the expectation values of P_{36}^{sfc} operator are close to zero.

In the deuteron case, $\langle P_{36}^{sfc} \rangle = -1/81$, it is a typical case of the second class. We collect some systems which have the same symmetry characteristics as deuteron in this subsection. The binding energies, E_b , and corresponding RMS , $\mathcal{R}$, of these systems are tabulated in Table 2.

Table 2. Binding energy, E_b , and corresponding RMS , $\mathcal{R}$, for the systems with $\langle P_{36}^{sfc} \rangle \approx 0$. The units for E_b and RMS are in MeV and fm , respectively.

S	System		$\langle P_{36}^{sf^c} \rangle$	Model I (Set I) $E_b // \mathcal{R}$	Model I (set II) $E_b // \mathcal{R}$
0	$NN(S=1, T=0)$ deuteron		-1/81	0.2 // 1.63	-0.7 // 1.68
-1	$N\Lambda$ ($S=0, T=1/2$)	$N\Lambda(S=0, T=1/2)$	0	-6.6 // 1.63	-7.7 // 1.69
		$N\Lambda - N\Sigma(S=0, T=1/2)$		-5.7 // 1.59	-7.2 // 1.66
	$N\Lambda$ ($S=1, T=1/2$)	$N\Lambda(S=1, T=1/2)$	0	-7.7 // 1.68	-7.7 // 1.68
		$N\Lambda - N\Sigma(S=1, T=1/2)$		-6.6 // 1.63	-6.7 // 1.64
		$N\Sigma(S=0, T=3/2)$	-1/81	-5.1 // 1.58	-5.3 // 1.59
-2	$\Lambda\Lambda$ ($S=0, T=0$)	$\Lambda\Lambda(S=0, T=0)$	0	-4.8 // 1.54	-5.6 // 1.58
		$\Lambda\Lambda - N\Sigma - \Sigma\Sigma(S=0, T=0)$			
		H particle		-2.0 // 1.15	8.2 // 0.91
-3		$\Lambda\Sigma(S=1, T=1/2)$	4/81	-8.1 // 1.74	-7.7 // 1.72
		$N\Omega(S=2, T=1/2)$	0	3.5 // 1.18	12.7 // 0.98
		$\Delta\Omega(S=3, T=3/2)$	0	4.4 // 1.15	14.2 // 0.96
-4		$\Xi\Xi(S=0, T=1)$	-1/81	4.1 // 1.17	0.4 // 1.30
-5	$\Xi\Omega$ ($S=1, T=1/2$)	$\Xi\Omega(S=1, T=1/2)$	1/81	9.5 // 1.02	4.2 // 1.14
		$\Xi\Omega - \Xi^*\Omega(S=1, T=1/2)$		32.9 // 0.78	32.6 // 0.77

The data in Table 2 shows that the deuteron is weakly bound in the Model I (Set I) case, which indicates that the chiral $SU(3)$ quark model, in principle, can reasonably describe the structure of deuteron². It is also seen that in the single $N\Lambda$ channel and the coupled $N\Lambda - N\Sigma$ channel with $S=0$ (or 1) and $T=1/2$ cases and the single $N\Sigma$ channel with $S=0$ and $T=3/2$ case, no matter which set of model parameters is employed, the systems are not bound. These are in agreement with experiments. The H particle is also not bound in Set I but weakly bound in Set II. The resultant mass of H is close to the $\Lambda\Lambda$ threshold in both Set I and Set II, and this feature is consistent with the recent finding in experiments [13]. These results further convince ourselves that the chiral $SU(3)$ quark model is reasonable and reliable in the bound-state study.

It should be noted that $\langle P_{36}^{sf^c} \rangle$ of these systems appoximately being zero means that the symmetry structure of this kind makes the quark-exchange effect less important, and consequently, the contribution from the kinetic energy term shows relatively repulsive nature to the kinetic energy of the relative motion between two well-separated interactive baryons (see Appendix), which makes interacting baryons apart. Therefore, very similar to deuteron, whether the two-baryon system is bound depends on the feature of the interaction between interacting baryons, especially that caused by chiral fields, namely the overall characteristics of the short- and medium-range $NPQCD$ effects, to a

² Of course, it is easy to fine-tune the mass of σ_0 so that the best agreement between the calculated and experimental binding energy of deuteron can be achieved. However, it is not necessary to give a very accurate mass of predicted dibaryon, we would relinquish this adjustment.

considerable extent. If the characteristics is attractive in nature, the system would be bound like deuteron, and we call it as a deuteron-like system. If it shows weak attraction or even repulsive feature, the system would not be bound anymore. It is also noticed that the OGE interaction provides repulsion in the deuteron, $N\Lambda$, $N\Sigma$, $\Xi\Xi(S=0, T=1)$ and $\Xi\Omega(S=1, T=1/2)$ systems. To form a bound state, a strong enough attractive interaction by the meson exchange must be requested. For instance, the π exchange, especially the tensor force, causes the weakly binding of deuteron and the relative strong attraction from the coupling between the chiral fields and quarks makes $\Xi\Xi(S=0, T=1)$ and $\Xi\Omega(S=1, T=1/2)$ bound. Then in the H particle case, although OGE interaction provides attraction, due to the repulsive effect from the kinetic energy term, the only attraction from OGE potential is not strong enough to cause binding. The attractive feature of the chiral field would be helpful to form a weakly bound H . But it is model parameter dependent. In the $N\Omega(S=2, T=1/2)$ and $\Delta\Omega(S=3, T=3/2)$ cases, OGE contributes nothing, the weakly bound behavior of these systems fully depends on the attractive features of the chiral fields. Moreover, it is noteworthy that very similar to the deuteron, the $\Lambda\Lambda - N\Xi - \Sigma\Sigma(S=0, T=0)$ and $\Xi\Omega - \Xi^*\Omega(S=1, T=1/2)$ systems are bounder than the $\Lambda\Lambda(S=0, T=0)$ and $\Xi\Omega(S=1, T=1/2)$ systems due to the channel coupling effect.

In summary, we predict that $N\Omega(S=2, T=1/2)$, $\Delta\Omega(S=3, T=3/2)$, $\Xi\Xi(S=0, T=1)$ and $\Xi\Omega(S=1, T=1/2)$ are weakly bound baryon-baryon states. The mass of the H particle is around the $\Lambda\Lambda$ threshold. $\Xi\Omega - \Xi^*\Omega(S=1, T=1/2)$ system is a bound state with a relative large binding energy due to the strong channel coupling.

3.2. The systems in which the expectation values of P_{36}^{sfc} operator are close to $-1/9$.

The systems in which $\langle P_{36}^{sfc} \rangle \approx -1/9$ are collected in this category. The binding energies, E_b , and corresponding root-mean-square radii, $\mathcal{R}$, of these systems are tabulated in Table 3.

Table 3. Binding energy, E_b , and corresponding RMS, $\mathcal{R}$, for the systems with $\langle P_{36}^{sfc} \rangle = -1/9$. The units for E_b and RMS are in MeV and fm , respectively.

$\mathcal{S}$	System	$\langle P_{36}^{sf^c} \rangle$	Model I (Set I) $E_b // \mathcal{R}$	Model I (set II) $E_b // \mathcal{R}$
0	$\Delta\Delta(S=3, T=0) (d^*)$	-1/9	22.2 // 1.01	18.5 // 1.05
	$\Delta\Delta(S=0, T=3)$	-1/9	16.0 // 1.10	13.5 // 1.14
-1	$\Sigma^*\Delta(S=0, T=5/2)$	-1/9	24.6 // 0.99	19.0 // 1.04
	$\Sigma^*\Delta(S=3, T=1/2)$	-1/9	25.9 // 0.95	29.3 // 0.93
-5	$\Xi^*\Omega(S=0, T=1/2)$	-1/9	92.4 // 0.71	76.5 // 0.72
-6	$\Omega\Omega(S=0, T=0)$	-1/9	116.1 // 0.66	98.5 // 0.67

From this table, one sees that all the systems in this category have the feature of $\langle P_{36}^{sf^c} \rangle = -1/9$. It indicates that the system has relatively higher anti-symmetry in the *spin-flavor-color* space. As a consequence, the contribution of the kinetic energy to the binding energy plays a relatively attractive role in comparison with the kinetic energy in two independent baryons [21] (see Appendix). This characteristic would bring six quarks closer. If the chiral fields can additionally provide attraction between interacting baryons, the deeply bound state may be established, such as $\Omega\Omega(S=0, T=0)$ and $\Xi^*\Omega(S=0, T=1/2)$. Even the contributions of chiral fields are mild, the system with the symmetry structure of this kind may still have the binding energy of several tens *MeV*. Therefore, in any case, the system in this category would be a bound state. However, it should be noticed that in most states here, such as $\Delta\Delta$ and $\Sigma^*\Delta$, both composed baryons have strong decay modes. Therefore, if the $\Delta\Delta$ and $\Sigma^*\Delta$ are not bound deeply enough, namely their masses are not smaller than the thresholds of $NN\pi\pi$ and $N\Lambda\pi\pi$, respectively, these four states should be bound states with broad widths. Only $\Omega\Omega(S=0, T=0)$ and $\Xi^*\Omega(S=0, T=1/2)$ are the most interesting systems. Both Ω 's in $\Omega\Omega(S=0, T=0)$ can decay only through the weak mode, so that $\Omega\Omega(S=0, T=0)$ is a bound state with a narrow width. It is also possible that the mass of $\Xi^*\Omega(S=0, T=1/2)$ is smaller than the threshold of $\Xi\Omega\pi$ (with Set I), then this state could also be a narrow width bound state.

4. Effects of interactions induced by chiral-quark field couplings.

In this section, we would demonstrate another factor which dominates the binding behavior of the system concerned, namely the interactions caused by the chiral-quark field couplings.

The characters of chiral fields are different case by case, and the importance and sensitivity of these fields in the bound-state problem of six-quark systems also dissimilar. In order to analyze the effect of the chiral field on the binding

energy of the baryon-baryon system, we build up other two models. In Model II, the π , K , η , η' and σ_0 are considered. This is so called extended chiral $SU(2)$ quark model. The Model III is called chiral $SU(2)$ quark model, in which only π and σ_0 fields are remained.

We arrange all concerned systems which are possibly bound into following types:

4.1. Deuteron-like systems.

To convince readers the the importance of the interaction for binding, we present the contributions of various terms, such as kinetic energy, OGE, pseudoscalar meson exchange and scalar mesons exchange terms to the total binding energy for the typical case $\Xi\Xi_{(0,1)}$ in table 4.

Table 4. Contributions of various terms to binding energy for $\Xi\Xi_{(0,1)}$, the unit for energy is in MeV .

	kine.	OGE	π	K	η	η'	σ	σ'	κ	ϵ
Model I Set I	-17.3	-15.2	-1.2	-1.7	-1.4	-0.8	37.7	1.9	-1.2	6.2
Model II	-12.0	-5.5	-0.8	-1.1	-0.8	-0.5	26.0	—	—	—
Model III	-16.1	-15.8	-1.4	—	—	—	40.5	—	—	—

From the table, we sees that although the kinetic energy, OGE and pseudoscalar mesons provide the repulsive feature, the strong attractive effect from scalar mesons, especially σ meson, would compensate the repulsion and make the $\Xi\Xi_{(0,1)}$ system weakly bound.

We then demonstrate the binding energies and the corresponding root-mean-square-radii of the bound systems in Sect.3.1 with Models I (set I), II and III, respectively, in Table 5, except the H particle which was carefully studied in our previous paper [5].

Table 5. Binding energy, E_b , and corresponding RMS, $\mathcal{R}$, for deuteron-like systems in various chiral quark models. The units for E_b and RMS are in MeV and fm , respectively. The system is denoted by symbol $[B_1B_2]_{(S,T)}$ with B_1 , B_2 , S and T being baryons 1 and 2 and the total spin and isospin of the system, respectively.

Model	$S = -5$	$S = -4$	$S = -3$		$S = 0$
	$[\Xi\Omega - \Xi^*\Omega]_{(1,1/2)}$	$[\Xi\Xi]_{(0,1)}$	$[N\Omega]_{(2,1/2)}$	$[\Delta\Omega]_{(3,3/2)}$	$d_{(S=1,T=0)}$
	$E_b // \mathcal{R}$	$E_b // \mathcal{R}$	$E_b // \mathcal{R}$	$E_b // \mathcal{R}$	$E_b // \mathcal{R}$
I (Set I)	32.9 // 0.78	4.1 // 1.17	3.5 // 1.18	4.4 // 1.15	0.2 // 1.63
II	29.3 // 0.79	-0.5 // 1.33	31.8 // 0.81	34.3 // 0.80	2.1 // 1.52
III	17.6 // 0.86	3.1 // 1.18	49.5 // 0.74	49.6 // 0.74	4.4 // 1.41

As mentioned in the preceding section, the binding energies of the systems in this type are sensitive to the contributions offered by the chiral field. This

character is very similar to that of deuteron and can explicitly be seen in this table. In most cases in this type, say the systems with low strange number, when one switches the model from Model II to Model I (Set I) or even from Model III to model II, namely the strange clouds are taken into account step by step, additional strange-cloud-caused interactions would make the overall inter-baryon interaction less attractive in nature, and consequently the systems would become less bounder. In a word, these systems are bounder in the chiral $SU(2)$ quark model than in the chiral $SU(3)$ quark model. But, in high strangeness systems, additionally taking strange clouds into account would benefit the binding.

The channel sketches and the predicted mass ranges of these states with the different combinations of chiral fields, which are denoted by shaded areas, are plotted in Figs.2(a)-(d) for the $[\Xi\Omega - \Xi^*\Omega]_{(1,1/2)}$, $[\Xi\Xi]_{(0,1)}$, $[N\Omega]_{(2,1/2)}$ and $[\Delta\Omega]_{(3,3/2)}$ systems, respectively.

4.2. $\Delta\Delta$ - and $\Omega\Omega$ -like systems

Now, we analyze the role or importance of the interaction in the system in the third class or in subsection 3.2.. As is mentioned above, except the symmetry character $\langle P_{36}^{sf_c} \rangle \approx -1/9$ of the system, the effect of the interaction is dominantly responsible for binding. A detailed study of the binding energies of the systems in this category show that, different with the contribution provided by the interaction in the second class, the contribution from exchange term of the interaction is substantially large and generally has two different character for various states in the class. Considering this difference together with the different decay mode, we further distinguish the states in this class into $\Delta\Delta$ -like states and $\Omega\Omega$ -like states. In $\Delta\Delta$ -like states, where only weak decays exist, the direct and exchange terms of interactions play comparable roles in forming dibaryon and the binding energies of these states are only a few tens of MeV. As a specific example, the calculated result shows that in $\Delta\Delta_{(0,3)}$, the exchange term of the interaction induced by σ meson provides a binding energy of 17.0 MeV, which is comparable to the direct contribution of 16.0 MeV. While in $\Omega\Omega$ -like states, where at most one strong decay mode exists, the exchange terms of interactions play much more important roles than the direct terms do and the binding energies of these states can reach nearly one

hundred MeV. For instance, in $\Omega\Omega_{(0,0)}$, the contribution of the exchange term of the σ interaction is 80.6 MeV, which is nearly two times larger than that of the direct contribution of 44.3 MeV. We would investigate these two types of states in detail in the following.

4.2.a. $\Delta\Delta$ -like systems.

We firstly single out $\Delta\Delta_{(0,3)}$ as a representative to decompose the binding energy of the state in this class. The results are tabulated in Table 6.

Table 6. Contributions of various terms to binding energy for $\Delta\Delta_{(0,3)}$, the unit for energy is in *MeV*.

		kine.	OGE	π	K	η	η'	σ	σ'	κ	ϵ
Model I	Set I	8.0	-26.7	-15.0	0.0	-2.3	-0.4	33.0	14.5	0.0	4.8
Model	II	6.2	-12.7	-10.0	0.0	-1.2	-0.2	24.1	—	—	—
Model	III	12.5	-17.8	-17.8	—	—	—	36.4	—	—	—

It is seen that only OGE and pseudosclar mesons provide repulsive contribution, while both kinetic energy and the scalar mesons devote their attractive contribution to binding. Moreover, it is shown that again the potential terms, especially the terms provided by the scalar meson, dominate the binding behavior.

The binding energies and the corresponding root-mean-square-radii for this type of states with Models I (set I), II and III are tabulated in Table 7, respectively.

Table 7. Binding energy, E_b , and corresponding *RMS*, $\mathcal{R}$, for $\Delta\Delta$ -like systems in various chiral quark models. The units for E_b and *RMS* are in *MeV* and *fm*, respectively. The system is denoted by symbol $[B_1 B_2]_{(S,T)}$ with B_1 , B_2 , S and T being baryons 1 and 2 and the total spin and isospin of the system, respectively.

Model	$S = -1$		$S = 0$	
	$[\Sigma^* \Delta]_{(0,5/2)}$ $E_b // \mathcal{R}$	$[\Sigma^* \Delta]_{(3,1/2)}$ $E_b // \mathcal{R}$	$[\Delta \Delta]_{(0,3)}$ $E_b // \mathcal{R}$	$[\Delta \Delta]_{(3,0)} (d^*)$ $E_b // \mathcal{R}$
I (Set I)	24.6 // 0.99	25.9 // 0.95	16.0 // 1.10	22.2 // 1.01
II	11.7 // 1.13	78.0 // 0.79	6.3 // 1.25	64.8 // 0.84
III	21.5 // 0.99	76.3 // 0.80	13.2 // 1.11	62.7 // 0.86

It is shown that the binding characters vary with the spin structures of systems. When one turns the model from Model I (Set I) to Model II, the binding energies of the $S = 0$ systems become smaller and those of the $S = 3$ systems become larger. However, if the model is switched from Model II to Model III, in comparison with the preceding case the binding energies for

different spin systems change in opposite directions. Nevertheless, substituting Model I (Set I) by Model III would cause the binding energy decreasing in the $S = 0$ system and increasing in the $S = 3$ system. Precisely, the systems with $S = 0$ and $S = 3$ are more bound in the chiral $SU(3)$ quark model and the extended chiral $SU(2)$ quark model, respectively. Moreover, because CMI shows repulsive feature in the $S = 0$ system and attractive nature in the $S = 3$ system, the binding energies of the $S = 3$ systems are generally higher than those of the $S = 0$ systems.

We also draw channel sketches and predicted mass ranges of $[\Sigma^*\Delta]_{(0,5/2)}$, $[\Sigma^*\Delta]_{(3,1/2)}$, $[\Delta\Delta]_{(0,3)}$, $[\Delta\Delta]_{(3,0)}$ states with different combination of chiral fields, which are signified by shaded areas, in Fig.2(e) and Fig.2(f), respectively. Since in the $\Sigma^*\Delta$ cases, the predicted masses are above the threshold of the strong decay channel $\Lambda N\pi\pi$, and in the $\Delta\Delta$ cases, the resultant masses are above the threshold of the strong decay channel $NN\pi\pi$, these bound states would have large widths. It might not be easy to detect in experiments.

It is worth to mention that because the Hamiltonian for the octet and decuplet baryons are in the same form and the calculated masses of these baryons can well-fit the data, the RGM calculations for the systems involving Δ are valid. Of course, whether this model treatment agrees with reality should be verified by the experiment in future.

4.2.b. $\Omega\Omega$ -like systems.

As what we did for deuteron- and $\Delta\Delta$ -like states, we choose $\Omega\Omega_{(0,0)}$ as a specific example to show the contributions of various Hamiltonian terms.

Table 8. Contributions of various terms to binding energy for $\Omega\Omega_{(0,0)}$, the unit for energy is in MeV .

	kine.	OGE	π	K	η	η'	σ	σ'	κ	ϵ
Model I Set 1	8.6	-82.7	0.0	0.0	-7.6	-15.0	124.9	0.0	0.0	87.7
Model II	12.3	-33.3	0.0	0.0	-6.5	-12.5	112.5	—	—	—
Model III	12.7	-58.5	0.0	—	—	—	97.7	—	—	—

The large binding energy of $\Omega\Omega_{(0,0)}$ in model I and the binding behaviors of the system in model II and III can clearly be understood by the detailed contributions of interactions in table 8. Apparently, the kinetic energy contribute attractively to the binding energy and the σ meson (in Model I, also ϵ meson) provides strong attractive contribution and dominates the binding behavior. OGE, η and η' are repulsive behavior contributors.

We tabulate the binding energies and the corresponding root-mean-square-radii for this type of systems with Models I (set I), II and III in Table 9, respectively.

Table 9. Binding energy, E_b , and corresponding RMS, $\mathcal{R}$, for $\Omega\Omega$ -like systems in various chiral quark models. The units for E_b and RMS are in MeV and fm , respectively. The system is denoted by symbol $[B_1 B_2]_{(S,T)}$ with B_1 , B_2 , S and T being baryons 1 and 2 and the total spin and isospin of the system, respectively.

Model	$S = -6$	$S = -5$
	$[\Omega\Omega]_{(0,0)}$	$[\Xi^*\Omega]_{(0,1/2)}$
	$E_b \quad // \mathcal{R}$	$E_b \quad // \mathcal{R}$
I (Set I)	116.1 // 0.66	92.4 // 0.71
II	74.2 // 0.69	58.8 // 0.75
III	53.5 // 0.73	52.1 // 0.76

It is seen that no matter in which model, these two states are bound states, or dibaryons. The *CM*I of the system in this type shows repulsive feature which offers disadvantage to binding. However, the benefit from the symmetry structure discussed in Sect.3.2 somehow overcomes this disadvantage. On the other hand, because these systems are high-strangeness systems, the attractive characteristics from the σ_0 - field-caused interaction and strange-cloud-induced interactions make the systems deeply bound and corresponding $\mathcal{R}$ s less than $0.76fm$. In other word, by using the chiral $SU(3)$ quark model, these systems could be even bound. More detailed analysis for $[\Omega\Omega]_{(0,0)}$ is given in Ref.[21].

The channel sketches and the predicted masses of the $[\Omega\Omega]_{(0,0)}$ and $[\Xi^*\Omega]_{(0,1/2)}$ states with different combinations of chiral fields, which are presented through shaded areas, are shown in Fig.2(g) and Fig.2(h), respectively. Since Ω does not have strong decay mode, the predicted $[\Omega\Omega]_{(0,0)}$ bound state should have narrow width. However, in the $[\Xi^*\Omega]_{(0,1/2)}$ system, because Ξ^* has the strong decay mode and the predicted masses of $\Xi^*\Omega$ by using different models stride over the threshold of the strong decay channel $\Xi\Omega\pi$, whether the width of the bound state is narrow depends on the model employed. With the chiral $SU(3)$ model, which, we believe, is a right model for the high strangeness system, the width should be narrow too.

Furthermore, we would emphasize that the purpose of this section is to trace the influences of various chiral fields on the calculated results through employing different models such as Models II and III. In our consideration, the

result with Model I is the most reliable one, because investigating a system with strange quarks without considering strange clouds is unreasonable.

5. Model parameter dependence

To ensure the reliability of our predictions, various kinds of model parameter-dependencies are also investigated.

It is clear that the variation of b_u would change the size of the baryon as well as the values of model parameters via the conditions mentioned in Sect.2, and the variation of m_s would change the strength of the OGE potential, g_s . We plot E_b with respect to b_u with different m_s for six most interesting systems, where $\langle P_{36} \rangle = -1/9$, in Fig.3. From this figure, one finds that the binding energy decreases with increasing values of b_u . When b_u is altered to a large value, the higher spin the system has, the milder the binding energy changes, while with increasing the mass of strange quark, the more strange quarks the system contains, the larger binding energy the system has. As long as the value stays in a reasonable region, say from $0.5fm$ to $0.55fm$, these systems remain the bound nature.

The m_σ -dependence is also studied, because the σ induced interaction is always a dominant piece that responses for the binding behavior for the two-baryon system. The results with different values of m_σ are tabulated in Table 7.

Table 10. The m_σ -dependence of the binding energy E_b . The units of E_b and m_σ are in MeV . The system is denoted by symbol $[B_1 B_2]_{(S,T)}$ with B_1 , B_2 , S and T being baryons 1 and 2 and the total spin and isospin of the system, respectively.

m_σ		550	600	625	650	700
E_b	$\Omega\Omega_{(0,0)}$	140.6	121.9	116.1	107.1	94.3
	$\Xi^*\Omega_{(0,1/2)}$	116.2	99.5	92.4	86.5	75.4
	$\Xi\Omega_{(1,1/2)}$	21.3	11.2	9.5	4.7	0.5
	$N\Omega_{(2,1/2)}$	17.7	7.1	3.5	0.9	-2.8
	$\Sigma^*\Delta_{(0,5/2)}$	36.5	28.1	24.6	21.9	17.0
	$\Sigma^*\Delta_{(3,1/2)}$	40.8	30.2	25.9	22.5	16.4
	$\Delta\Delta_{(0,3)}$	25.5	18.7	16.0	13.9	10.2
	$\Delta\Delta_{(3,0)}$	35.0	25.8	22.2	19.3	14.2
	$NN_{(1,0)}(d)$	5.1	1.6	0.2	-0.5	-2.1

From this table, one sees that as long as m_σ is in a reasonable region, namely the existent experimental data can unifiably and reasonably be explained, most systems studied would not change their binding behaviors, ex-

cept those whose binding energies are marginal, for instance, when $m_\sigma = 700\text{MeV}$, $N\Omega$ becomes unbound.

The cutoff mass dependence is also investigated. Our result shows that if one increases all the cutoff masses to 1.2GeV , the binding energies of the systems concerned would slightly arisen. The binding behaviors are relative stable.

In a word, as long as the model parameters are selected in reasonable regions and all the experimental data, such as NN and NY scattering data, the ground state masses of baryons, binding features of well-known baryon-baryon states, and etc., can be basically reproduced, the concerned six two-baryon systems with $\langle P_{36}^{sf_c} \rangle = -1/9$ should be bound, especially, $\Omega\Omega_{(0,0)}$ and $\Xi^*\Omega_{(0,1/2)}$ should be deeply bound dibaryons.

6. Concluding remarks.

The possible candidates of S -wave baryon-baryon bound states with various strange numbers are systematically studied by using the chiral quark model. In terms of the chiral $SU(3)$ quark model, which are believed to be one of the best models in the quark-gluon degrees of freedom in well reproducing as much experimental data as possible, we predict the two-baryon bound states without any additional parameters.

There are two major factors that affect the binding behavior of the two-baryon system. One is the symmetry property of the system, which is characterized by the matrix-element $\langle \mathcal{A}_{36}^{sf_c} \rangle$, and the other is the interactions between quarks including both the direct and exchange components, especially the interactions caused by chiral-quark field couplings.

We classify two-baryon bound states (even almost bound states) into three types. The first type state is the deuteron-like state. The symmetry structure of the system in this type shows that the quark exchange effect is not so important, namely, there is no serious Pauli blocking effect. If the chiral field can provide enough attraction between interacting baryons, the bound state can be formed. This kind of bound state might also be obtained by using the models on baryon level. For instance, in terms of Nijmegen model, one also

predicted some bound states in the $\mathcal{S} = -2, -3$ and -4 systems [27].

The $\Delta\Delta$ -like state is the second type state. The system of this type is mostly symmetric in the *orbit* space. In these states, the interaction, especially the scalar meson induced interaction, dominates the binding behavior. Together with the symmetry behavior of the system, the relatively deeply-bound state can be formed. However, because both baryons in the system have strong decay modes, only when the predicted binding energy is lower than the threshold of the strong decay channel, say the $NN\pi\pi$ channel for the $\Delta\Delta$ system and the $\Lambda N\pi\pi$ channel for the $\Sigma^*\Delta$ system, the width of the bound state could be narrow. In our calculation, the predicted binding energies of $\Delta\Delta$ and $\Sigma^*\Delta$ are not large enough, so that the widths of these bound states should be rather broad. Although this kind of bound state might not easily be detected in experiments, it may worth to search in the future.

The third type state is the $\Omega\Omega$ -like state, which is the most interesting state in our study. Same as those in the second type, the system in this type is mostly symmetric in the *orbit* space. Meanwhile, the strange chiral fields can offer rather strong attraction. As a result, these states are deeply bound states. If we believe that the chiral $SU(3)$ quark model is one of the most suitable models in describing the system with high strangeness, the predicted binding energy of $\Xi^*\Omega$ is possibly below the threshold of the strong decay channel $\Xi\Omega\pi$. Thus, both $\Omega\Omega$ and $\Xi^*\Omega$ can merely have weak decays, and consequently are deeply bound states with narrow widths.

It should be specially emphasized that the states of the second and third types possess the six-quark structure, and their inter-baryon distances are relatively short. These characteristics cannot be provided by the model on the baryon level. Thus, they are new dibaryon systems. The existences of this kind of dibaryons would be an important place to reveal *QCD* phenomenology.

Appendix.

In the framework of RGM [28], the upper bound is given by the expectation

value of the Hamiltonian

$$\langle H \rangle = \frac{\langle \Psi | H | \Psi \rangle}{\langle \Psi | \Psi \rangle}, \quad (24)$$

where $H = T + V$ is the Hamiltonian, with T and V being the kinetic energy operator and the potential operator, respectively, Ψ represents the trial wave function of the system and $\langle H \rangle$ denotes the upper bound of the system. The kinetic operator of a six-quark system can be written as:

$$T = \sum_{i=1}^6 T_i - T_{CM}, \quad (25)$$

where T_i and T_{CM} denote the kinetic energy operators of the i -th quark and of the center of mass motion (CM), respectively. Substituting the trial wave function (Eq.(21)) and the antisymmetrizer $\mathcal{A}'$ (Eq.(22)) into projection equation

$$\langle [\hat{\phi}_A \hat{\phi}_B \chi_{CM}(\vec{R}_{CM})]_{ST} | H - E | \mathcal{A}' [\hat{\phi}_A \hat{\phi}_B \chi_{rel}(\vec{R}) \chi_{CM}(\vec{R}_{CM})]_{ST} \rangle = 0, \quad (26)$$

one obtains RGM equation

$$\begin{aligned} \{ & - \frac{\hbar^2}{2\mu} \nabla_{\vec{R}}^2 + V_{rel}^{dir}(\vec{R}) - E_{rel} \} \chi_{rel}(\vec{R}) \\ & + \int [K^T(\vec{R}, \vec{R}') + K^V(\vec{R}, \vec{R}') - E_{tot} N^{exch}(\vec{R}, \vec{R}')] \chi_{rel}(\vec{R}') d\vec{R}' = 0, \end{aligned} \quad (27)$$

where $V_{rel}^{dir}(\vec{R})$ represents the relative potential between two clusters, E_{tot} and E_{rel} denote the total energy and the relative energy between two clusters, respectively, and $K^T(\vec{R}, \vec{R}')$, $K^V(\vec{R}, \vec{R}')$ and $N^{exch}(\vec{R}, \vec{R}')$ describe the kinetic energy exchange kernel, the potential energy exchange kernel and the normalization exchange kernel, respectively. Expanding $\chi_{rel}(\vec{R})$ by well-defined basis functions

$$\chi_{rel}(\vec{R}) = \sum_{i=1}^n c_i \phi_{rel}(\vec{R}, \vec{S}_i), \quad (28)$$

and left-multiplying $\phi_{rel}(\vec{R}, \vec{S}_i)$ to RGM equation and integrating over $\vec{R}$ and $\vec{R}'$, one obtains the secular equation of the bound state problem,

$$\begin{aligned} & \sum_{i=1}^n [H_{ij} - E_{tot} N_{ij}] c_j \\ & = \sum_{i=1}^n [H'_{ij} - E_{rel} N_{ij}] c_j = 0, \quad (i = 1, \dots, n) \end{aligned} \quad (29)$$

with H_{ij} and N_{ij} being the Hamiltonian and normalization kernels, respectively, and

$$E_{tot} = E_{in} + E_{rel}, \quad (30)$$

E_{tot} , E_{in} and E_{rel} being the total, inner cluster and relative energies, respectively. Apparently, H' can be written as:

$$\begin{aligned} H'_{ij} &= H_{ij} - E_{in}N_{ij} \\ &= [T_{ij} - E_{in}^T N_{ij}] + [V_{ij} - E_{in}^V N_{ij}]. \end{aligned} \quad (31)$$

In this equation, T_{ij} and V_{ij} is the kinetic energy and potential energy kernels, respectively, and the superscripts T and V denote the kinetic energy and potential energy parts, respectively. Then,

$$\begin{aligned} H_{ij}^{T'} &= T_{ij} - E_{in}^T N_{ij} \\ &= (T_{rel}^{dir})_{ij} + (T_{rel}^{exch})_{ij} \end{aligned} \quad (32)$$

with

$$(T_{rel}^{dir})_{ij} = \langle \phi_{rel}(\vec{R}'', \vec{S}_i) | (-\frac{\hbar^2}{2\mu} \nabla_{\vec{R}'}^2) \delta(\vec{R}'' - \vec{R}') | \phi_{rel}(\vec{R}', \vec{S}_j) \rangle \quad (33)$$

and

$$(T_{rel}^{exch})_{ij} = \langle \phi_{rel}(\vec{R}'', \vec{S}_i) | K^T(\vec{R}'', \vec{R}') | \phi_{rel}(\vec{R}', \vec{S}_j) \rangle. \quad (34)$$

To compute $H_{ij}^{T'}$ explicitly, we employ the Generating Coordinate Method (GCM) technique, which is equivalent to RGM. In GCM, we re-write

$$\Psi = \sum_{i=1}^n c_i \psi_i \quad (35)$$

with

$$\begin{aligned} \psi_i &= \mathcal{A}'[\hat{\phi}_A \hat{\phi}_B \phi_{rel}(\vec{R}', \vec{S}_i) \chi_{CM}(\vec{R}_{CM}) \chi_{ST} \chi_c] \\ &= \mathcal{A}'[\prod_{n=1}^3 \phi_{0s}(\vec{r}_n - \frac{\vec{S}_i}{2}) \prod_{k=4}^6 \phi_{0s}(\vec{r}_k + \frac{\vec{S}_i}{2}) \chi_{ST} \chi_c] \end{aligned} \quad (36)$$

and

$$\phi_{0s}(\vec{r}_n - \frac{\vec{S}_i}{2}) = N_n \exp -\frac{1}{2b_n^2} (\vec{r}_n - \frac{\vec{S}_i}{2})^2. \quad (37)$$

To ensure the total CM motion can explicitly be separated, we keep ω unchanged, namely

$$\omega = (m_u b_u^2)^{-1} = (m_s b_s^2)^{-1}. \quad (38)$$

If one temporarily ignores the flavor symmetry breaking, namely $b = b_u = b_s$ and $m = m_u = m_s$, for instance in the $(\Omega\Omega)$ or $\Delta\Delta$ or NN system, the following discussion would be flavor independent and it is easy to obtain the kinetic energy induced S-wave adiabatic *effective interaction*

$$\begin{aligned} E_{ii}^T &= \frac{H_{ii}^{T'}}{N_{ii}} - \frac{K_0}{3} \\ &= 2K_0 x \left[-1 + Ctgh(3x) \frac{1 - 9\langle P_{36}^{sfc} \rangle Cosh(x)/Cosh(3x)}{1 - 27\langle P_{36}^{sfc} \rangle Sinh(x)/Sinh(3x)} \right] \end{aligned} \quad (39)$$

with

$$x = \frac{S_i^2}{4b^2} \quad (40)$$

and

$$K_0 = \frac{3}{4}\hbar\omega. \quad (41)$$

According to the definition of the binding energy

$$E_b = -(M_{DB} - M_A - M_B), \quad (42)$$

where M_{DB} , M_A and M_B denote the masses of the dibaryon, baryon A and baryon B, respectively, the contribution of the kinetic energy to the binding energy of the dibaryon in the adiabatic approximation should be

$$E_b^T = -\sum_{i=1}^n c_i^2 E_b^T(S_i) \quad (43)$$

with

$$E_b^T(S_i) = E_{ii}^T + \frac{K_0}{3}, \quad (44)$$

where c_i is the wavefunction. We plot $E_b^T(S_i)$ versus S_i in Fig.4 (for convenience, we drop the subscript i in S_i in Fig.4). In this figure, the solid and

dashed curves denote the results with $\langle P_{36}^{sfc} \rangle = -1/9$ and $\langle P_{36}^{sfc} \rangle = -1/81$, respectively. Apparently, when $\langle P_{36}^{sfc} \rangle = -1/81$, $E_b^T(S_i)$ has less attractive character than that in the $\langle P_{36}^{sfc} \rangle = -1/9$ case. And when $\langle P_{36}^{sfc} \rangle \geq 0$, it would be even less attractive.

Apparently, E_b^T is always a negative number (or repulsive for binding). This indicates that, when off-diagonal matrix elements contribution can be neglected under adiabatic approximation, E_b^T always has a repulsive effect.

However, when calculating E_b^T , one should also take the off-diagonal matrix elements into account. Further calculation shows that the contribution of the off-diagonal matrix elements differs substantially in the system with $\langle P_{36}^{sfc} \rangle = -1/9$ to that in the system with $\langle P_{36}^{sfc} \rangle \approx 0$. In the former case, it is so large that E_b^T can be changed to a positive value (or attractive for binding). For instance, in the $\Omega\Omega_{(0,0)}$ dibaryon, the contribution of the diagonal matrix elements is -145.6 MeV while the contribution of the off-diagonal matrix elements is 154.2 MeV, and consequently $E_b^T = 8.6$ MeV. On the other hand, in the latter case, this contribution is not large enough to change the sign of E_b^T . For instance, in the deuteron case, the contribution of the diagonal matrix elements is -57.5 MeV while the contribution of the off-diagonal matrix elements is 43.0 MeV, and consequently $E_b^T = -14.5$ MeV. The effect of the off-diagonal matrix elements can also simply be seen by diagonalizing the $-(E_{ij}^T + \frac{K_0}{3})$, called $\tilde{E}_{ij}^T$, matrix. In the $\langle P_{36}^{sfc} \rangle = -1/9$ case, some of diagonalized matrix elements $\tilde{E}_{ii}^T$ can be positive. Because

$$E_b^T = \sum_i^n \tilde{c}_i^2 \tilde{E}_{ii}^T, \quad (45)$$

where $\tilde{E}_{ii}^T$ is the diagonalized matrix element, and $\tilde{c}_i$ is the wave function after the unitary transformation that diagonalizes the $-(E_{ij}^T + \frac{K_0}{3})$ matrix, the resultant E_b^T becomes positive. While for the system with $\langle P_{36}^{sfc} \rangle \approx 0$, the diagonalized matrix elements are always negative, consequently, E_b^T always keeps a negative value.

The physics picture of the positive contribution (or attractive contribution) of the kinetic energy operator to the total binding energy may be understood as the following. In the dibaryon with $\langle P_{36}^{sfc} \rangle = -1/9$, such as $\Omega\Omega_{(0,0)}$, six quarks interfere with each other very strongly due to the large quark exchange effect,

so that they cannot move as freely as they do in two independent baryons. So the expectation value of kinetic energy operator in such system (which is definitely positive) would be smaller than the kinetic energy in two independent baryons. While in system with $\langle P_{36}^{sc} \rangle \approx 0$, the quark exchange effect is so weaker that the expectation value of kinetic energy operator is always larger than that the kinetic energy in two baryons.

References

- [1] R.L.Jaffee, *Phys. Rev. Lett.* **38**, 195 (1977).
- [2] A.Chodos, R.L.Jaffee, K.Johnson, et al., *Phys. Rev. D* **9**, 3471 (1974); T.DeGrand, R.L.Jaffee, K.Johnson, et al., *Phys. Rev. D* **12**, 2060 (1975).
- [3] M.Oka and K.Yazaki, *Phys. Lett. B* **90**, 41 (1980); A.Faessler, F.Fernandez, G.Lübeck, et al., *Nucl. Phys. A* **402**, 555 (1983); M.Oka, K.Shimizu, K.Yazaki, *Nucl. Phys. A* **464**, 700 (1987); U.Straub, Z.Y.Zhang, K.B.Bräuer, A.Faessler, *Phys. Lett. B* **200**, 241 (1988); *Nucl. Phys. A* **483**, 686 (1988); A.Valcarce, B.Buchmann, A.Fernandez, A.Faessler, *Phys. Rev. C* **50**, 2246 (1994).
- [4] K.Yazaki, *Prog.Theor.Phys. Suppl.* **91**, 146 (1987), and references therein.
- [5] P.N.Shen, Z.Y.Zhang, Y.W.Yu, S.Q.Yuan, S.Yang, *J. Phys. G* **25**, 1807 (1999); *Chin. Phys. Lett.* , (1999).
- [6] K.Shimizu, M.Koyama, *Nucl. Phys. A* **646**, 211 (1999).
- [7] Z.Y.Zhang, Y.W.Yu, P.N.Shen, L.R.Dai, A.Faessler, U.Straub, *Nucl. Phys. A* **625**, 59 (1997).
- [8] S.Yang, P.N.Shen, Z.Y.Zhang, Y.W.Yu, *Nucl. Phys. A* **635**, 146 (1998).
- [9] G.S. Adkins et al., *Nucl. Phys. B* **228**, 552 (1983); E.M. Nyman, *Phys. Lett. B* **142**, 388 (1984).
- [10] H.Georgi, *Weak Interactions and Modern Particle Theory* (Benjamin/Cummings)
- [11] I.T.Obukhovskiy and V.G.Kusainov, *Phys. Lett. B* **238**, 142 (1990); A.Buchmann, E.Fernandez and K.Yazaki, *Phys. Lett. B* **269**, 35 (1991); E.M.Henley and G.A.Miller, *Phys. Lett. B* **251**, 453 (1991).
- [12] M.D.Scadron, *Phys. Rev. D* **26**, 239 (1982).
- [13] K.Imai, **Hadrons and Nuclei with Strangeness- Proceedings of the First Sino-Japan Symposium on Strangeness Physics**, (to be published in 1999), and references therein.

- [14] Y.W. Yu, Z.Y.Zhang and X.Q. Yuan, *High Energy Phys. and Nucl. Phys.* **23**, 859 (1999).
- [15] M.Oka, K.Shimizu, K.Yazaki, *Phys. Lett. B* **130**, 365 (1983); K.Yazaki, **Hadrons and Nuclei with Strangeness- Proceedings of the First Sino-Japan Symposium on Strangeness Physics**, (to be published in 1999), and references therein..
- [16] Y.W.Yu, Z.Y.Zhang and X.Q.Yuan, *Communi.Theor.Phys.* **31**, 1 (1999).
- [17] D.Kamae and T.Fujita, *Phys. Rev. Lett.* **38**, 471 (1977).
- [18] N.R.Walet, *Phys. Rev. C* **48**, 2222 (1993).
- [19] F.Wang, J.L.Ping, G.H.Wu, et al., *Phys. Rev. C* **51**, 3411 (1995)
- [20] X.Q.Yuan, Z.Y.Zhang, Y.W.Yu and P.N.Shen, *Phys. Rev. C* **60**, 045203 (1999); *Communi.Theor.Phys.* **32**, 169 (1999).
- [21] Z.Y.Zhang, Y.W.Yu, C.R. Ching, et al., *Phys. Rev. C* **61**, 065204 (2000), and references therein. P.N.Shen, Q.B.Li, Z.Y.Zhang, et al., Proceedings of Hadron'99, Beijing, China, 1999, *Nucl. Phys. A*.
- [22] R.Bilger and H.A.Clement, *Phys. Rev. Lett.* **71**, 42 (1993).
- [23] G.Wagner, L.Ya.Glozman, A.J.Buchmann and A.Faessler, *Nucl. Phys. A* **594**, 263 (1995), and references therein.
- [24] Q.B.Li and P.N.Shen, *Chin. Phys. Lett.* **17**, 182 (2000).
- [25] Particle Data Group, *Euro.Phys.J. C* **3**, 353 (1998).
- [26] P.J.G.Mulders, A.T.Aerts and J.J.Swarts, *Phys. Rev. Lett.* **40**, 1543 (1978); *Phys. Rev. D* **21**, 2653 (1980).
- [27] T.A.Rijken, *Proceedings of Asia Pasific Conference on Few-Body Problems In Physics*, Noda/Kashiwa, Japan, 1999.
- [28] K.Wildermuth and Y.C.Tang, *A Unified Theory of the Nucleus*, Academic Press Inc., 1977.

Figure captions

Fig.1. Λ - P scattering.

Fig.2. Energy level sketches for a. $[\Xi\Omega-\Xi^*\Omega]_{(1,1/2)}$, b. $[\Xi\Xi]_{(0,1)}$, c. $[N\Omega]_{(2,1/2)}$, d. $[\Delta\Omega]_{(3,3/2)}$, e. $[\Sigma^*\Delta]_{(0,5/2)}$ and $[\Sigma^*\Delta]_{(3,1/2)}$, f. $[\Delta\Delta]_{(0,3)}$ and $[\Delta\Delta]_{(3,0)}$, g. $[\Omega\Omega]_{(0,0)}$, h. $[\Xi^*\Omega]_{(0,1/2)}$.

Fig.3. Binding energy E_b with respect to b_u and m_s for systems with $\langle\mathcal{A}^{sc}\rangle \approx 2$. The solid curves are for $m_s=470$ MeV, the dashed curves are for $m_s=515$ MeV.

Fig.4. In the S-wave case, the contribution to binding energy provided by the kinetic-energy related terms with different S in the adiabatic approximation.







